

GENERATION GAPS AND ABELIANISED DEFECTS OF FREE PRODUCTS

KARL W. GRUENBERG AND PETER A. LINNELL

ABSTRACT. Let G be a group of the form $G_1 * \cdots * G_n$, the free product of n subgroups, and let M be a $\mathbb{Z}G$ -module of the form $\bigoplus_{i=1}^n M_i \otimes_{\mathbb{Z}G_i} \mathbb{Z}G$. We shall give formulae in various situations for $d_{\mathbb{Z}G}(M)$, the minimum number of elements required to generate M . In particular if C_1, C_2 are nontrivial finite cyclic groups of coprime orders, $G = (C_1 \times \mathbb{Z}) * (C_2 \times \mathbb{Z})$ and $F/R \cong G$ is the free presentation obtained from the natural free presentations of the two factors, then the number of generators of the relation module, $d_{\mathbb{Z}G}(R/R')$, is three. It seems plausible that the minimum number of relators of G should be 4, and this would give a finitely presented group with positive relation gap. However we cannot prove this last statement.

1. INTRODUCTION

Given a group G and a $\mathbb{Z}G$ -module M , let $d(G)$ denote the minimal number of generators of G , let $d_G(M)$ denote the minimal number of $\mathbb{Z}G$ -module generators of M , and let ΔG denote the augmentation ideal of G , that is the ideal of $\mathbb{Z}G$ generated by $\{g - 1 \mid g \in G\}$. Then the *generation gap* of the finitely generated group G is the difference

$$\text{gap } G := d(G) - d_G(\Delta G).$$

If G is finitely presentable and $R \hookrightarrow F \twoheadrightarrow G$ is a finite free presentation (meaning F is a finitely generated free group and R has a finite number of generators as a normal subgroup of F), then the *abelianised defect* $\text{adef}(F/R)$ of the free presentation is the difference $d_G(R/R') - d(F)$, where R' is the commutator group of R and the free abelian group R/R' is viewed as a $\mathbb{Z}G$ -module (the relation module of the presentation). A fair amount is known about the numerical functions gap and adef for finite groups. Our purpose here is to make a start with their study for infinite groups. In particular, to indicate how results on finite groups can be used to discuss free products. In the next section, after reviewing some relevant facts of the finite theory, we state and discuss our main results. We mention just two striking consequences:

1.1. Proposition. *If G is the free product $G_1 * \cdots * G_n$ of the finite groups $G_1, \dots, G_n$ of mutually coprime orders, then $\text{gap}(G) = 0$ if and only if there exists a positive integer k such that $\text{gap}(G_k) = 0$ and G_i is cyclic for all $i \neq k$.*

Date: Mon Dec 24 16:37:04 EST 2007.

2000 Mathematics Subject Classification. Primary: 20F05; Secondary: 20E06.

Key words and phrases. generation gap, relation gap, free product.

The first author died on 10 October 2007. The second author gratefully acknowledges financial support from the EPSRC.

1.2. Proposition. *Let G_1, G_2 be nontrivial finite cyclic groups of coprime orders, let C_1, C_2 be infinite cyclic groups, and let $G = (G_1 \times C_1) * (G_2 \times C_2)$. If $F/R \simeq G$ is the free presentation obtained from the natural 2-generator free presentations of the two factors, then $d_G(R/R') = 3$.*

The second of these results could have an important consequence for the relation gap problem: given a finite free presentation $R \hookrightarrow F \twoheadrightarrow G$ of a group G , then the *relation gap* of the presentation is defined to be $d_F(R) - d_G(R/R')$, where $d_F(R)$ denotes the minimum number of elements needed to generate R as a normal subgroup of F . No examples are known of positive relation gap. The group of Proposition 1.2 above needs at most 4 (obvious) relations. If 4 really equals $d_F(R)$, then F/R would have positive relation gap. Proposition 1.2 was mentioned in the interesting survey article [10] by Jens Harlander.

A similar result was proved in [2, Proposition 1], where consequences to the “ $D(2)$ conjecture” [13] were discussed. We shall show how to obtain [2, Propositions 1,2] from Theorem 4.6 at the end of Section 6.

The first author Karl Gruenberg died on 10 October 2007. At that time this paper was more or less complete; mistakes that remain are entirely the fault of the second author, Peter Linnell. Karl was a close and generous friend to not only the second author, but many other mathematicians. His influence through his research and teaching will be greatly missed.

2. REVIEW AND OVERVIEW

Among finite groups, generation gap zero is the rule rather than the exception: all simple groups (indeed all 2-generator groups) and all soluble groups have generation gap zero. However there do exist groups of arbitrarily large generation gap. The smallest group with nonzero generation gap is known to be the direct product of 20 copies of the alternating group of degree 5 [5]. For more on these matters, see [7, §13].

Finite groups also behave well with respect to abelianised defect: on a given finite group G , the function $\text{adef}(F/R)$ as F/R varies over all finite free presentations of G is constant (cf. [7, §14]). Hence $\text{adef}(G)$ is an unambiguous notation for this integer. For finitely presentable groups G , the set of $\text{adef}(F/R)$ for varying finite free presentations F/R is bounded below; so the lower bound may be written $\text{adef}(G)$. A free presentation $F/R \simeq G$ is called *minimal* if $d(F) = d(G)$.

If $G = G_1 * \cdots * G_n$ and we are given free presentations $F_i/R_i \simeq G_i$ for $i = 1, \dots, n$, then $F = *F_i$ projects onto G with kernel the normal closure of $R_1, \dots, R_n$.

If R is a relation group, we shall always abbreviate the resulting relation module R/R' as $\overline{R}$. When the relation group is R_i , we shall use the streamlined notation $\overline{R}_i$ for $\overline{R}_i$.

2.1. Theorem. *Let $G = G_1 * \cdots * G_n$ with all the G_i finite groups of mutually coprime orders. Then*

(a)

$$d_G(\Delta G) = \max_k \{d_{G_k}(\Delta G_k)\} + n - 1$$

whence

$$\text{gap}(G) = \min_k \{ \text{gap}(G_k) + \sum_{i \neq k} (d(G_i) - 1) \}.$$

- (b) Let $F/R \simeq G$ be the free presentation constructed from free presentations $F_i/R_i \simeq G_i$ of the free factors. Then

$$d_G(\overline{R}) = \max_k \{d_{G_k}(\overline{R}_k) + \sum_{i \neq k} d(F_i)\},$$

whence $\text{adef}(F/R) = \max_k \{\text{adef}(G_k)\}$.

An immediate consequence of Theorem 2.1(a) is Proposition 1.1. Theorem 2.1(b) tells us, for example, that if each G_i is an elementary abelian p_i -group of rank 2 and $d(F_i) = 2$, then $d_G(\overline{R}) = 3 + 2(n-1) = 2n+1$ because here (clearly) $d_{G_i}(\overline{R}_i) = 3$. One might expect $d_F(R)$ to be $3n$ since R is the normal closure of $R_1, \dots, R_n$ and $d_{F_i}(R_i) = 3$ for all i . However in the case $n = 2$, Hog-Angeloni, Lustig and Metzler showed that $d_F(R) = 5$ [11, Addendum to Theorem 3 on p. 163] and hence the relation gap of F/R is zero. Their results yield for general n that $d_F(R) \leq 3n-1$. If there exists $n > 2$ such that $d_F(R) = 3n-1$, then the group involved will have positive relation gap, namely $3n-1 - (2n+1) = n-2 > 0$.

Theorem 2.1 follows from the general Theorem 4.6 stated and proved in §4. All our results on gap and adef are consequences of this. Theorem 4.6 is a join of two results, the first of which, Theorem 4.4, is our major contribution: it gives an upper bound for the minimal number of generators of a $\mathbb{Z}G$ -module M , where G has subgroups $G_1, \dots, G_n$ and M contains “good” $\mathbb{Z}G_i$ -submodules M_i that together G -generate M . When $G = G_1 * \dots * G_n$ and $M = \bigoplus_{i=1}^n (M_i \otimes_{\mathbb{Z}G_i} \mathbb{Z}G)$, our upper bound for $d_G(M)$ becomes a lower bound by a remarkable theorem of George Bergman [1]. The restriction to what we call “good” modules (cf. Definition 4.3) is probably unavoidable. Fortunately it is no restriction at all for finite groups. The proof of this has no bearing on any of our other arguments and we therefore postpone it to an appendix.

The following two theorems look somewhat like Theorem 2.1, but here the factors are infinite groups and this leads to difficulties with the “good” property of modules.

2.2. Theorem. *Let $H = H_1 * \dots * H_n$, where each $H_i = G_i \times A_i$, G_i is a finite nilpotent group, A_i is a finitely generated nilpotent group with A/A' torsion free and the orders of $G_1, \dots, G_n$ are mutually coprime. Then*

(a)

$$d_H(\Delta H) = \max_k \{d(H_k) + \sum_{i \neq k} (d(A_i) + \delta_{A_i,1})\},$$

where $\delta_{A_i,1} = 1$ when $A_i = 1$ (equivalently $d(A_i) = 0$) and $\delta_{A_i,1} = 0$, when $A_i \neq 1$.

(b) $\text{gap}(H) = \min_k \{\sum_{i \neq k} (d(G_i) - \delta_{A_i,1})\}$.

We conclude that a free product of finitely generated abelian groups, with torsion groups of mutually coprime orders, has generation gap zero if, and only if, every factor but one is torsion free or finite cyclic. Theorem 2.2 is an immediate corollary of the more general Theorem 6.5 in §6.

Finally, we state the theorem of which Proposition 1.2 is a special case.

2.3. Theorem. *Let $H = H_1 * \dots * H_n$, where each $H_i = G_i \times C_i$ with G_i finite cyclic and C_i infinite cyclic. Assume that the orders of $G_1, \dots, G_n$ are mutually coprime. Take natural minimal free presentations $F_i/R_i \simeq H_i$ and let $F/R \simeq H$ be the resulting free presentation of H . Then $d_H(\overline{R}) = n+1$.*

Theorem 2.3 is an immediate corollary of Theorem 6.13. It seems probable to us that Theorem 2.3 remains true (in an obviously modified enunciation) when each C_i is allowed to be free abelian of finite rank and G_i is a suitably restricted finite group. Just what may be needed to establish such a generalisation will become clear in §6.

3. PRELIMINARIES

We explain some notation and collect a number of results, mostly well-known, that are needed later. Modules will be right modules and mappings will be written on the left. If V is a finitely generated module over a ring S , we write $d_S(V)$ for the minimum number of elements needed to generate V as an S -module. Of course if $X \subseteq V$, then XS indicates the S -submodule of V generated by X . In the case $S = kG$, the group ring of the group G over the commutative ring k , we usually write $d_G(V)$ instead of $d_{kG}(V)$. If G is finite, we let $\hat{G} = \sum_{g \in G} g \in kG$. If K is a commutative ring containing k , then KV means $K \otimes_k V$, so if V is a $\mathbb{Z}G$ -module, $d_G(\mathbb{Q}V)$ means $d_{\mathbb{Q}G}(\mathbb{Q} \otimes_{\mathbb{Z}} V)$. For a prime p , we shall let $\mathbb{F}_p$ indicate the field with p elements. The commutator subgroup of G will be denoted by G' and we let $G^n = \langle g^n \mid g \in G \rangle$ for any positive integer n . If H is a group, then $\delta_{G,H} \in \mathbb{Z}$ is the Kronecker δ : so $\delta_{G,H} = 1$ if $G \cong H$ and $\delta_{G,H} = 0$ if $G \not\cong H$. Thus the δ in Theorem 2.2 is the Kronecker delta.

If π is a set of primes, then π' denotes the complementary set. Let $\pi(G)$ denote the set of primes dividing the orders of the finite subgroups of G . If G is torsion free, set $\pi(G) = \emptyset$. Let M be a finitely generated $\mathbb{Z}G$ -module. We call M a *Swan module* if there exists a prime p such that $d_G(M) = d_G(M/pM)$; in that case we call p an *M -prime*. If G is finite and M is a $\mathbb{Z}G$ -lattice, then M -primes, if they exist, are divisors of $|G|$ (cf. 3.3). There is a well developed theory of Swan $\mathbb{Z}G$ -lattices when G is finite: cf. [9, §7.1] for the basics. Augmentation ideals and relation modules of finite groups are Swan lattices. A *relation prime* of G is an $\overline{R}$ -prime, where $\overline{R}$ is the relation module coming from a finite free presentation of G . When G is finite, this notion is independent of the chosen presentation: if $\overline{R}_1, \overline{R}_2$ are relation modules, then the $\overline{R}_1$ -primes coincide with the $\overline{R}_2$ -primes: cf. [9, Lemma 7.13].

3.1. Lemma. *Let H be a finitely generated group and let G be a homomorphic image of H such that $d(G) = d(H)$. If $\text{gap}(G) = 0$ and ΔG is a Swan module for G , then $\text{gap}(H) = 0$ and ΔH is a Swan module for H . Moreover every ΔG -prime is a ΔH -prime.*

Proof. If p is a ΔG -prime, then $d(H) = d(G) = d_G(\Delta G) = d_G(\Delta G/p\Delta G) \leq d_H(\Delta H/p\Delta H) \leq d_H(\Delta H) \leq d(H)$. $\square$

3.2. Corollary. *If H is a finitely generated nilpotent group, then $\text{gap}(H) = 0$ and ΔH is a Swan module. When H/H' is torsion free, every prime is a ΔH -prime; otherwise every ΔH -prime belongs to $\pi(H/H')$.*

Proof. There exists a prime p such that $G := H/H'H^p$ satisfies $d(G) = d(H)$. Since G is an elementary abelian p -group, it has generation gap zero. When H/H' is torsion free, any prime will do for p ; but if H/H' has torsion subgroup $T \neq 1$, then p must divide $|T|$. $\square$

3.3. Lemma. *Let G be a finite group and let M be a finitely generated $\mathbb{Z}G$ -module. Then $d_G(\mathbb{Q}M) \leq d_G(M/pM)$ for every prime p , while $d_G(\mathbb{Q}M) = d_G(M/pM)$ for every $p \notin \pi(G) \cup \pi(M)$.*

Proof. First we show that $d_G(\mathbb{Q}M) \leq d_G(M/Mp)$. Choose $x_1, \dots, x_n \in M$ to generate M as a $\mathbb{Z}G$ -module modulo p . Then the elements $x_i g$ for $1 \leq i \leq n$ and $g \in G$ generate M as a $\mathbb{Z}$ -module modulo p . Since M is finitely generated as a $\mathbb{Z}$ -module, we see that the elements $x_i g$ generate $\mathbb{Q}M$ as a $\mathbb{Q}$ -module. Hence the elements x_i generate $\mathbb{Q}M$ as a $\mathbb{Q}G$ -module.

Next we show that $d_G(M/Mp) \leq d_G(\mathbb{Q}M)$ when $p \notin \pi(G) \cup \pi(M)$; this will complete the proof. Choose $x_1, \dots, x_n \in M$ to generate $\mathbb{Q}M$ as a $\mathbb{Q}G$ -module. Write $N = x_1 \mathbb{Z}G + \dots + x_n \mathbb{Z}G$. Then $\mathbb{Q}N = \mathbb{Q}M$, consequently $\mathbb{C} \otimes_{\mathbb{Z}} N \cong \mathbb{C} \otimes_{\mathbb{Z}} M$. If M' is the torsion group of M and $\overline{M} = M/M'$, then $\overline{M}$ is a $\mathbb{Z}G$ -lattice and $\overline{M}/p\overline{M} \cong M/pM$ since p is prime to $|M'|$. Let k denote the algebraic closure of the field $\mathbb{F}_p$. By Brauer character theory, we see by looking at the decomposition numbers that $k \otimes_{\mathbb{Z}} M$ and $k \otimes_{\mathbb{Z}} N$ have the same composition factors as kG -modules. Since p is prime to $|G|$, all kG -modules are completely reducible and we deduce that $k \otimes_{\mathbb{Z}} M \cong k \otimes_{\mathbb{Z}} N$. Therefore $M/Mp \cong N/Np$ as $\mathbb{F}_p G$ -modules by the Noether-Deuring theorem. Thus $d_G(M/pM) \leq n$ as required. $\square$

4. THE MAIN TECHNICAL RESULT

Given a finitely generated $\mathbb{Z}G$ -module M and a finite set of primes π , let $(X_p)_{p \in \pi}$ be a family of subsets of M such that X_p is a minimal $\mathbb{Z}G$ -generating set of M modulo pM and $X_p \supseteq X_q$ if, and only if, $d_G(M/pM) \geq d_G(M/qM)$. We call $(X_p)_{p \in \pi}$ a *nested π -family in M* . Note that the nested structure ensures that if X_q is the minimal element of $(X_p)_{p \in \pi}$, then for every $p \in \pi$,

$$(4.1) \quad d_G(M/(pM + X_q \mathbb{Z}G)) = d_G(M/pM) - d_G(M/qM).$$

If $\pi = \{p_1, \dots, p_t\}$, we will write $X_i = X_{p_i}$ for $1 \leq i \leq t$ and assume that $X_1, \subseteq \dots, \subseteq X_t$.

4.2. Lemma. *For every M and π there exists a nested π -family in M .*

Proof. Let $\pi = \{p_1, \dots, p_t\}$ be ordered so that $d_G(M/p_i M) \geq d_G(M/p_j M)$ if, and only if, $i \leq j$. Choose a subset Y_i of M so that its image in $M/p_i M$ is a minimal $\mathbb{Z}G$ -generating set of $M/p_i M$. Then $|Y_1| \geq |Y_2| \geq \dots \geq |Y_t|$.

Let $d = |Y_1|$ and enumerate the elements in each Y_i using zeros if necessary at the end: $Y_i = \{y_{i,1}, y_{i,2}, \dots, y_{i,d}\}$. Let $p'_i = \prod_{j \neq i} p_j$. Then $(p_i, p'_i) = 1$ and $p_i m_i + p'_i m'_i = 1$ for suitable integers m_i, m'_i . We define

$$x_k = \sum_{i=1}^t p'_i m'_i y_{i,k} \quad (1 \leq k \leq d).$$

Thus $x_k \equiv p'_j m'_j y_{j,k} \equiv y_{j,k}$ modulo $p_j M$, whence $X_i = \{x_1, \dots, x_{d_i}\}$ with $d_i = |Y_i|$ gives the required nested π -family $X_1 \supseteq X_2 \supseteq \dots \supseteq X_t$. $\square$

4.3. Definition. Let G be a group and let M be a finitely generated $\mathbb{Z}G$ -module. We say M is *good* for G if there exists a finite set of primes $\pi(G, M)$ such that

- (g1) $d_G(M/pM)$ has constant value $\delta_G(M)$ for all $p \notin \pi(G, M)$, while $\delta_G(M) \leq d_G(M/pM)$ for all $p \in \pi(G, M)$;

(g2) for every finite set of primes π properly containing $\pi(G, M)$ there exists a nested π -family whose minimal element X satisfies

$$d_G(M/(pM + X\mathbb{Z}G)) \leq 1 \quad \text{for all } p \notin \pi,$$

and $M/X\mathbb{Z}G$ is a torsion group of finite exponent.

It is then automatic that $|X| = \delta_G(M)$. Also if $p \in \pi'$, then $d_G(M/(pM + X\mathbb{Z}G)) = 1$ if, and only if, $p \in \pi(M/X\mathbb{Z}G)$ (because $M \neq pM + X\mathbb{Z}G$ if, and only if, $p \in \pi(M/X\mathbb{Z}G)$).

4.4. Theorem. *Let G be a group containing subgroups $G_1, \dots, G_n$, let M_i be a good $\mathbb{Z}G_i$ -module and let M be a $\mathbb{Z}G$ -module containing $M_1, \dots, M_n$ and generated by them as a $\mathbb{Z}G$ -module. Assume that either all the M_i are Swan modules or that $d_{G_i}(M_i/pM_i) > \delta_{G_i}(M_i)$ for some integer i and some prime p . Then*

$$d_G(M) \leq \max_p \left\{ \sum_{i=1}^n d_{G_i}(M_i/pM_i) \right\}.$$

Proof. Set $\delta = \max_p \left\{ \sum_{i=1}^n d_{G_i}(M_i/pM_i) \right\}$, $e_i = \delta_{G_i}(M_i)$ and let π be a finite set of primes strictly containing $\pi(G_1, M_1) \cup \dots \cup \pi(G_n, M_n)$. Choose a nested π -family in M_1 as required for (g2) and let X_1 be the minimal element. Set $N_1 = M_1/X_1\mathbb{Z}G_1$. Now $\pi(N_1)$ is finite and $d_{G_1}(N_1/pN_1) \neq 0$ if, and only if, $p \in \pi(N_1)$. Define $\pi_2 = \pi \cup \pi(N_1)$ and choose a nested π_2 -family in M_2 as required for (g2). Let its minimal element be X_2 and set $N_2 = M_2/X_2\mathbb{Z}G_2$. Continue in this manner, producing $X_3, \dots, X_n$, with $N_i = M_i/X_i\mathbb{Z}G_i$ and $\pi_i = \pi \cup \pi(N_1) \cup \dots \cup \pi(N_{i-1})$ for $i = 2, \dots, n$. This has ensured that if $i < j$ and $p \in \pi'$, then $d_{G_i}(N_i/pN_i)$ and $d_{G_j}(N_j/pN_j)$ cannot both be non-zero. For suppose $d_{G_i}(N_i/pN_i) \neq 0$. Then $p \in \pi(N_i)$, whence $p \in \pi_j$ and then, since X_j is the minimal element in a nested π_j -family, $d_{G_j}(N_j/pN_j) = d_{G_j}(M_j/pM_j) - \delta_{G_j}(M_j)$ by (4.1), where the right hand side is zero by (g1), because $p \notin \pi(G_j, M_j)$. If $d_{G_j}(N_j/pN_j) \neq 0$, then $p \notin \pi_j$ (otherwise $d_{G_j}(N_j/pN_j) = 0$ as we have just seen) and so $p \in \pi(N_i)'$, which ensures $d_{G_i}(N_i/pN_i) = 0$. Thus we have

$$(4.5) \quad \sum_{i=1}^n d_{G_i}(N_i/pN_i) \leq 1 \quad \text{for every } p \in \pi'.$$

Suppose $\delta = \sum_{i=1}^n e_i$. Since $e_i \leq d_{G_i}(M_i/pM_i)$ for every p by (g1), we see that $e_i = d_{G_i}(M_i/pM_i)$ for every p and every i . In this case our hypothesis makes every M_i a Swan module, and so $e_i = d_{G_i}(M_i)$ by using an M_i -prime p . Thus $\delta = \sum_{i=1}^n d_{G_i}(M_i)$, which gives the result. Therefore we may assume that $\delta > \sum_{i=1}^n e_i$.

For each positive integer $r \leq \delta - \sum_{i=1}^n e_i$, let $\mathcal{P}_r$ denote the set of primes $p \in \pi$ for which $r \leq \sum_{i=1}^n d_{G_i}(N_i/pN_i)$. Thus $\mathcal{P}_1 \neq \emptyset$ (because $e_i < d_{G_i}(M_i/pM_i)$ for some i and some $p \in \pi$), $\mathcal{P}_1 \supseteq \mathcal{P}_2 \supseteq \dots \supseteq \mathcal{P}_s$, where $s = \delta - \sum_{i=1}^n e_i$ and $\mathcal{P}_{s+1} = \emptyset$.

Recall $N_i \neq pN_i$ for some $p \in \pi'$ if and only if $\pi(N_i) \cap \pi' \neq \emptyset$. Let $\pi_0(N_i) = \pi(N_i) \cap \pi'$ and let I_0 be the set of all i such that $\pi_0(N_i) \neq \emptyset$. By (4.5), the sets $\pi_0(N_i)$, for $i \in I_0$, are disjoint. If $i \in I_0$, we know $d_{G_i}(N_i/pN_i) = 1$ for every $p \in \pi_0(N_i)$ (property (g2)), whence we may choose a p -element $u_{i,0}(p)$ which $\mathbb{Z}G_i$ -generates N_i modulo pN_i . If $i \notin I_0$ set $u_{i,0}(p) = 0$ for all p .

We now define inductively $\mathbb{Z}G_i$ -quotients $N_{i,k}$ of N_i for $1 \leq k \leq s$, starting with $N_{i,1} = N_i$, for $1 \leq i \leq n$.

The set of primes $p \in \pi$ so that $N_i \neq pN_i$ for some i is precisely $\mathcal{P}_1$. Let $\pi_1(N_i) = \pi(N_i) \cap \pi$ and let I_1 be all i so that $\pi_1(N_i) \neq \emptyset$. With $i \in I_1$, choose a p -element $u_{i,1}(p) \in N_i$ so that $d_{G_i}(N_i/(pN_i + u_{i,1}(p)\mathbb{Z}G_i)) < d_{G_i}(N_i/pN_i)$. If $\pi_1(N_i) = \emptyset$, put $u_{i,1}(p) = 0$ for all p . Let $\pi(N_{i,1})$ be the union of the disjoint sets $\pi_0(N_i)$, $\pi_1(N_i)$ and define

$$K_{i,1} = \{u_{i,0}(p), u_{i,1}(p) \mid p \in \pi(N_{i,1})\}\mathbb{Z}G_i.$$

Set $N_{i,2} = N_i/K_{i,1}$.

If $N_{i,2} \neq pN_{i,2}$, then $p \in \mathcal{P}_2$ (because then $N_i \neq pN_i$, $p \in \pi_1(N_i)$ and $d_{G_i}(N_i/pN_i) \geq 2$). Thus $\pi(N_{i,2}) \subseteq \mathcal{P}_2$. Let I_2 be all i for which $\pi(N_{i,2}) \neq \emptyset$. If $p \in \pi(N_{i,2})$, choose a p -element $u_{i,2}(p) \in N_i$ so that $d_{G_i}(N_{i,2}/(pN_{i,2} + \bar{u}_{i,2}(p)\mathbb{Z}G_i)) < d_{G_i}(N_{i,2}/pN_{i,2})$, where $\bar{u}_{i,2}(p) = u_{i,2}(p) + pN_{i,2}$. If $\pi(N_{i,2}) = \emptyset$, set $u_{i,2}(p) = 0$ for all p . Define

$$K_{i,2} = K_{i,1} + \{u_{i,2}(p) \mid p \in \pi(N_{i,2})\}\mathbb{Z}G_i$$

and $N_{i,3} = N_i/K_{i,2}$. Continue until we reach $N_i = K_{i,r}$, for some $r \leq s$ (because $\pi(N_{i,k}) \subseteq \mathcal{P}_k$ for all k).

By construction, for each i and each p , the set of all non-zero elements $u_{i,k}(p)$ for $k \geq 0$ maps to a minimal set of $\mathbb{F}_p G_i$ -generators of N_i/pN_i . When $u_{i,0}(p) \neq 0$, the set consists just of this one element because then $p \notin \pi$. The number of non-zero $u_{i,k}(p)$ for all $k \geq 0$ and all i is $\sum_{i=1}^n d_{G_i}(N_i/pN_i)$. Thus

$$\max_p |\{u_{i,k}(p) \mid k \geq 0 \text{ and all } i\}| \leq s.$$

Let $N = \bigoplus_{i=1}^n N_i \otimes_{\mathbb{Z}G_i} \mathbb{Z}G$. Now $A = \langle u_{i,k}(p) \mid \text{all } i, k, p \rangle$ is a *finite* additive subgroup of N and $N = A\mathbb{Z}G$ (because $N_i = K_{i,s}$ for all i). The subset $\{u_{i,k}(p) \mid \text{all } i, k\}$ generates $A(p)$, the Sylow p -subgroup of A and $d(A) = \max_p d(A(p))$. Hence $d_G(N) \leq \max_p d(A(p)) \leq s$.

Thus finally, $d_G(M) \leq d_G(N) + |X_1| + \cdots + |X_n| \leq s + \sum e_i = \delta$. $\square$

All our results on free products are consequences of the following theorem which is an immediate consequence of Theorem 4.4 and a theorem of Bergman [1, Theorem 2.3]:

4.6. Theorem. *Let $G = G_1 * \cdots * G_n$, let M_i be a good $\mathbb{Z}G_i$ -module for $1 \leq i \leq n$, and let $M = \bigoplus_{i=1}^n M_i \otimes_{\mathbb{Z}G_i} \mathbb{Z}G$. If either all the M_i are Swan modules or $d_{G_i}(M_i/pM_i) > \delta_{G_i}(M_i)$ for some i and some prime p , then*

$$d_G(M) = \max_p \left\{ \sum_{i=1}^n d_{G_i}(M_i/pM_i) \right\}.$$

Proof. Theorem 4.4 gives the inequality $\leq$, while Bergman's theorem tells us that, for each p , we have $\sum_{i=1}^n d_{G_i}(M_i/pM_i) = d_G(M/pM)$. $\square$

Note that, by Bergman's theorem, the conclusion of Theorem 4.6 is equivalent to the statement that M is a Swan $\mathbb{Z}G$ -module.

5. FINITE FREE FACTORS

If G is a finite group, then every finitely generated $\mathbb{Z}G$ -module M is good for G if we take $\pi(G, M) = \pi(G) \cup \pi(M)$. We already know that (g1) holds: this is Lemma 3.3 with $\delta_G(M) = d_G(\mathbb{Q}M)$. We postpone the proof of (g2) to the Appendix.

For the rest of this section we assume that $G_1, \dots, G_n$ are finite groups and $G = G_1 * \cdots * G_n$. We consider various applications of Theorem 4.6, beginning with

augmentation ideals. Here $\bigoplus_{i=1}^n \Delta G_i \otimes_{\mathbb{Z}G_i} \mathbb{Z}G = \Delta G$ (e.g. [8, §8.6]), each ΔG_i is a Swan lattice and $d_{G_i}(\mathbb{Q}\Delta G_i) = 1$ (which makes (g2) a triviality). Thus Theorem 4.6 gives

$$(5.1) \quad d_G(\Delta G) = \max_p \left\{ \sum_{i=1}^n d_{G_i}(\Delta G_i / p\Delta G_i) \right\}.$$

Hence *the augmentation ideal of a free product of finite groups is a Swan module*. When $G_1, \dots, G_n$ have mutually coprime orders, then (5.1) with Lemma 3.3 gives Theorem 2.1(a).

Next consider relation modules. Let $R_i \hookrightarrow F_i \twoheadrightarrow G_i$ be a free presentation and $R \hookrightarrow F \twoheadrightarrow G$ the resulting free presentation of G . Then $\overline{R} = \bigoplus_{i=1}^n \overline{R}_i \otimes_{\mathbb{Z}G_i} \mathbb{Z}G$ and $d_{G_i}(\mathbb{Q}\overline{R}_i) = d(F_i)$ (see e.g. [9, Theorem 2.7]). From Theorem 4.6 we obtain

$$(5.2) \quad d_G(\overline{R}) = \max_p \left\{ \sum_{i=1}^n d_{G_i}(\overline{R}_i / p\overline{R}_i) \right\}.$$

This has Theorem 2.1(b) as corollary. Furthermore, the Gruško-Neumann theorem tells us that a free presentation of the free product of the groups G_i is of the form $*_i F_i \twoheadrightarrow *_i G_i$ with each F_i a free group mapping onto G_i , consequently (5.2) shows that *the relation module of a finite free presentation of a free product of finite groups is a Swan module*.

The augmentation ideal of a group G may be viewed as the first kernel of a $\mathbb{Z}G$ -free resolution of $\mathbb{Z}$, while a relation module is the second kernel of such a resolution. We indicate how Theorem 2.1 may be generalised to arbitrary kernels.

Let H be a finite group, let

$$(5.3) \quad \dots \longrightarrow (\mathbb{Z}H)^{e_2} \xrightarrow{\phi_2} (\mathbb{Z}H)^{e_1} \xrightarrow{\phi_1} \Delta H \xrightarrow{\phi_0} 0$$

be a $\mathbb{Z}H$ -free resolution of ΔH and write $K_i = \ker \phi_i$. We recall a criterion for a lattice to be a Swan module. This is originally due to Swan, but we use the improved form due to Jacobinski [12].

Let $\mathbb{Z}_{(H)}$ denote the semilocal subring $\{a/b \mid a, b \in \mathbb{Z}, (b, |H|) = 1\}$ of $\mathbb{Q}$ and take a minimal free presentation $K \hookrightarrow (\mathbb{Z}_{(H)}H)^d \twoheadrightarrow (K_s)_{(H)}$, where $(K_s)_{(H)} = K_s \otimes_{\mathbb{Z}(H)}$. Then K_s is a Swan lattice if $\mathbb{Q}K$ contains a copy of every non-trivial irreducible $\mathbb{Q}H$ -module. Since H is a finite group, every finitely generated $\mathbb{Q}H$ -module is uniquely a direct sum of irreducible $\mathbb{Q}H$ -modules and we have

$$\mathbb{Q}K - (d - e_s + e_{s-1} - \dots + (-1)^s e_1) \mathbb{Q}H + (-1)^s \mathbb{Q}\Delta H = 0$$

which shows that $\mathbb{Q}\Delta H$ is a summand of $\mathbb{Q}K$ whenever s is odd and also when s is even provided $d - e_s + e_{s-1} - \dots + (-1)^s e_1 \neq 1$; but if the sum is 1, then $K \cong \mathbb{Z}_{(H)}$ and $\mathbb{Z}$ has projective period ℓ with $s + 2 \equiv 0 \pmod{\ell}$.

Finally $\mathbb{Q}K_s = (\mathbb{Q}H)^e \oplus (-1)^s \mathbb{Q}\Delta H$, where $e = e_s - e_{s-1} + \dots + (-1)^{s-1} e_1$. Thus $\mathbb{Q}K_s$ needs $e + 1$ $\mathbb{Q}H$ -module generators if s is even and e generators when s is odd. If K_s fails to be a Swan lattice, $\max_p \{d_H(K_s / pK_s)\} = d = e + 1 = d_H(\mathbb{Q}K_s)$. Thus $d_H(K_s / pK_s)$ has constant value for all primes p .

Now return to $G = G_1 * \dots * G_n$. Let

$$(5.4) \quad \dots \longrightarrow (\mathbb{Z}G_i)^{f_{i,2}} \xrightarrow{\theta_{i,2}} (\mathbb{Z}G_i)^{f_{i,1}} \xrightarrow{\theta_{i,1}} \Delta G_i \xrightarrow{\theta_{i,0}} 0$$

be a G_i -free resolution of ΔG_i . Applying $\otimes_{\mathbb{Z}G_i} \mathbb{Z}G$ for each i , we obtain a $\mathbb{Z}G$ -free resolution of $\Delta G_i \otimes_{\mathbb{Z}G_i} \mathbb{Z}G$. The sum of these resolutions yields a G -free resolution of ΔG :

$$(5.5) \quad \dots \longrightarrow (\mathbb{Z}G)^{f_2} \xrightarrow{\theta_2} (\mathbb{Z}G)^{f_1} \xrightarrow{\theta_1} \Delta G \xrightarrow{\theta_0} 0,$$

where $f_s = f_{1,s} + \dots + f_{n,s}$ and $\ker \theta_s = \bigoplus_{i=1}^n \ker \theta_{i,s} \otimes_{\mathbb{Z}G_i} \mathbb{Z}G$.

5.6. Proposition. *Let $G = G_1 * \dots * G_n$ and take resolutions (5.4) and (5.5). Suppose s is a non-negative integer such that if G_i has cohomological period ℓ_i , then $s + 2 \not\equiv 0 \pmod{\ell_i}$. Then*

$$(a) \quad d_G(\ker \theta_s) = \max_p \left\{ \sum_{i=1}^n d_{G_i}(\ker \theta_{i,s}/p \ker \theta_{i,s}) \right\};$$

(b) *if $G_1, \dots, G_n$ have mutually coprime orders, then*

$$d_G(\ker \theta_s) = \max_k \{ d_{G_k}(\ker \theta_{k,s}) + \sum_{i \neq k} d_{G_i}(\mathbb{Q} \ker \theta_{i,s}) \}$$

and $d_{G_i}(\mathbb{Q} \ker \theta_{i,s}) = f_{i,s} - f_{i,s-1} + \dots + (-1)^{s-1} f_{i,1} + \delta_s$, where $\delta_s = 1$ or 0 according as s is even or odd.

Proof. By our discussion of H above, $\ker \theta_{i,s}$ is a Swan lattice if s is as stated. Therefore Theorem 4.6 yields (a); then (b) is a consequence of (a), again using facts proved about H . $\square$

Another application of Theorem 4.6 in the context of finite groups concerns finitely generated nilpotent groups. If H is such a group, then there is a prime p so that the elementary abelian image $H/H' H^p$ has the same minimal number of generators as does H . Let us call such a prime a *generating prime* for H .

5.7. Proposition. *Let $H = H_1 * \dots * H_n$, where each H_i is a finitely generated nilpotent group. If every H_i , except possibly one, is cyclic or has H_i/H'_i torsion-free, then $\text{gap}(H) = 0$.*

Proof. Suppose (without loss of generality) that H_1 is neither cyclic nor has H_1/H'_1 torsion-free. Let q be a generating prime for H_1 and $J(q)$ be the set of all j such that q is a generating prime for H_j . This absorbs all H_j with H_j/H'_j torsion-free.

Pick a generating prime p_i for each i , but take $p_i = q$ whenever $i \in J(q)$. Set $G_i = H_i/H'_i H^{p_i}$. Then $G = G_1 * \dots * G_n$ is a homomorphic image of H and $d(G) = d(H)$. By reordering if necessary we may assume $J(q) = \{1, \dots, r\}$. Then G_i is cyclic for all $i > r$ and $p_i \neq q$. Hence

$$\max_p \left\{ \sum_{i=1}^n d_{G_i}(\Delta G_i/p \Delta G_i) \right\} = \sum_{i=1}^r d_{G_i}(\Delta G_i/q \Delta G_i) + n - r = d(G)$$

and so $\text{gap } G = 0$ (by Theorem 4.6) whence $\text{gap } H = 0$ by Lemma 3.1. $\square$

It would be interesting to have a converse of Proposition 5.7. The closest we come in this direction is Theorem 2.2 (cf. Theorem 6.5).

6. INFINITE FREE FACTORS

In order to use Theorem 4.6 for infinite groups, we must find good Swan modules. We begin with augmentation modules.

6.1. Proposition. *Let A be a non-trivial finitely generated nilpotent group with A/A' torsion-free and let G be a finite group such that*

- (a) $d(G) = d(G/G')$ and
- (b) $d_G(\Delta G/p\Delta G) = d(G/G'G^p)$ for all $p \in \pi(G)$.

If $H = G \times A$, then $\text{gap } H = 0$, ΔH is both a Swan lattice and a good $\mathbb{Z}H$ -module for $\pi(H, \Delta H) = \pi(G)$. Furthermore $d(H) = d(G) + d(A)$ and $\delta_H(\Delta H) = d(A)$.

For the proof of Proposition 6.1, we require the following result.

6.2. Lemma. *Let G be a finite group, let B be a finite nilpotent group and let $E = G \times B$. For any prime p ,*

$$d_E(\Delta E/p\Delta E) = \max\{d_G(\Delta G/p\Delta G), \dim G/G'G^p + \dim B/B'B^p\}.$$

Proof. If $B(p)$ is the Sylow p -subgroup of B , $d_B(\Delta B/p\Delta B) = d(B(p))$ ([9, Lemma 7.17(i)]) and $d(B(p)) = d(B/B'B^p)$. The result now follows from [3, Theorem 2p]. $\square$

Lemma 6.2 shows that every finite nilpotent group G has property (b) of Proposition 6.1. There are plenty of non-nilpotent finite groups having properties (a) and (b). Let $G = P \times B$, where $P = P'$, $B' = 1$, $\pi(P) \subseteq \pi(B)$ and $d(B/B^p) \geq d_P(\Delta P)$ for all $p \in \pi(G)$. For any $k \geq 1$, the direct power $G^{(k)}$ of G satisfies $d_{G^{(k)}}(\Delta G^{(k)}/p\Delta G^{(k)}) = k d(B/B^p)$ for all $p \in \pi(G)$ (again use [3, Theorem 2p]). Thus $G^{(k)}$ has property (b) for all $k \geq 1$. If k is large enough, then by a theorem of Wiegold (e.g. [9, Theorem 6.13]) $G^{(k)} = k d(G/G') = d(G^{(k)}/G^{(k)'})$, whence $G^{(k)}$ has property (a).

Proof of Proposition 6.1. When $G = 1$ the result follows from Corollary 3.2, so henceforth we assume $G \neq 1$. Set $\bar{A} = A/A^{|G|}$ and $\bar{H} = G \times \bar{A}$. Now $\text{gap}(G/G') = 0$ implies $\text{gap}(G) = 0$, by 3.1, and this implies $\text{gap}(\bar{H}) = 0$ (use [6, (2.3)] and note that what is called in that paper the presentation rank $\text{pr}(E)$ of the finite group E coincides with $\text{gap}(E)$: [7, §12]). It will follow from Lemma 3.1 that $\text{gap}(H) = 0$ and that ΔH is a Swan module, provided that $d(H) = d(\bar{H})$.

If $n \geq 2$ is a positive integer, C is a finite abelian group such that $nC = 0$, and D is a finitely generated free abelian group, then $d(C \times D/nD) = d(C) + d(D)$. Applying this with $n = |G|$, $C = G/G'$ and $D = A/A'$, we find that

$$d(H) \leq d(G) + d(A) = d(G/G') + d(A) = d((G/G') \times (\bar{A}/\bar{A}')) \leq d(\bar{H}) \leq d(H).$$

Thus $d(H) = d(\bar{H})$ and also $d(H) = d(G) + d(A)$.

Consider condition (g1). Let $a_1, \dots, a_e$ be free generators of A . By Corollary 3.2, $\{a_1 - 1, \dots, a_e - 1\}$ is a minimal set of $\mathbb{Z}A$ -module generators of ΔA modulo $p\Delta A$ for every prime p . Choose $p \notin \pi(G)$. Since $d_G(\Delta G/p\Delta G) = 1$ (by Lemma 3.3) there exists $x \in \Delta G$ so that $x\mathbb{Z}G + p\Delta G = \Delta G$. Let

$$(6.3) \quad X = \{x + (a_1 - 1)\hat{G}, a_2 - 1, \dots, a_e - 1\}$$

(where $\hat{G}$ is defined in the first paragraph of §3). We claim $X\mathbb{Z}H + p\Delta H = \Delta H$. For if $z \mapsto \bar{z}: \Delta H \twoheadrightarrow \Delta H/p\Delta H$ denotes the natural epimorphism and $W = \bar{X}\mathbb{Z}H$, then

$$(6.4) \quad (x + \hat{G}(a_1 - 1))(|G| - \hat{G}) = |G|x$$

shows $\bar{x} \in W$, whence $\hat{G}(\bar{a}_1 - 1) \in W$. Also $\bar{g} - 1 \in W$ for all $g \in G$ since $\bar{x}\mathbb{Z}H$ contains $(\Delta G + p\Delta H)/p\Delta H$. Hence $|G| - \hat{G} + p\Delta H \in W$, which gives

$$|G|(\bar{a}_1 - 1) = (|G| - \hat{G} + \hat{G})(\bar{a}_1 - 1) \in W,$$

and so $(\bar{a}_1 - 1) \in W$. Thus $W = \Delta H/p\Delta H$. We conclude $d_H(\Delta H/p\Delta H) \leq e$ and hence equality because $\Delta H \twoheadrightarrow \Delta A$. Thus $\delta_H(\Delta H) = e$ and of course $d_H(\Delta H/p\Delta H) \geq e$ for every prime p , whence property (g1).

Finally property (g2). Choose a finite set of primes π properly containing $\pi(G)$ and find a nested π -family $Y_1 \supseteq Y_2 \supseteq \dots \supseteq Y_t$ in ΔG that satisfies (g2). We may assume that there is a positive integer $s \leq t$ such that $p_i \in \pi(G)$ for $i < s$ and $p_i \notin \pi(G)$ for $i \geq s$. Let $Y_i = \{y_1, \dots, y_{d_i}\}$ (so $Y_t = \{y_1\}$). Define

$$X_t = \{y_1 + (a_1 - 1)\hat{G}, a_2 - 1, \dots, a_e - 1\}$$

for $i \geq s$, and for $i < s$ define

$$X_i = \{y_1 + (a_1 - 1)\hat{G}, a_2 - 1, \dots, a_e - 1, a_1 - 1, y_2, \dots, y_{d_i}\}.$$

Then $X_1 \supseteq \dots \supseteq X_t$ and we claim this is a nested π -family in ΔH . First consider $i < s$. Since X_i generates the same $\mathbb{Z}H$ -submodule as $S := \{y_1, y_2, \dots, y_{d_i}, a_1 - 1, a_2 - 1, \dots, a_e - 1\}$ and clearly $S\mathbb{Z}H + p_i\Delta H = \Delta H$, we need to show that $d_H(\Delta H/p_i\Delta H) = d_i + e$. With $\bar{H} = G \times \bar{A}$ as above, $H \twoheadrightarrow \bar{H}$ induces a bijection $S \simeq \bar{S}$ (since the kernel of $\Delta H \twoheadrightarrow \Delta \bar{H}$ is $(\Delta A^{|\hat{G}|})\mathbb{Z}H$). So we are reduced to proving $d_{\bar{H}}(\Delta \bar{H}/p_i\Delta \bar{H}) = d_i + e$. This is true by property (b) on G and Lemma 6.2.

Next consider the case $i \geq s$. Since X_i is the set X of (6.3) with $x = y_1$ and $p = p_i$, we see that X_t minimally generates ΔH modulo $p_i\Delta H$. Simplify the notation now by setting $X = X_i (= X_t)$ and $x = y_1$. Let q be a prime not in π . Then (by (g2) for ΔG), $d_G(\Delta G/(q\Delta G + x\mathbb{Z}G)) \leq 1$, whence there exists $z \in \Delta G$ such that $\{x, z\}\mathbb{Z}G + q\Delta G = \Delta G$. Then $X \cup \{z\}$ generates ΔH modulo $q\Delta H$.

To finish the proof, we must show that $\Delta H/X\mathbb{Z}H$ has finite exponent. Suppose m is the exponent of the finite module $\Delta G/x\mathbb{Z}G$. Then m is prime to p ($= p_i$) and (6.4) shows that $|G|x \in X\mathbb{Z}H$. So $|G|m\Delta G$ is contained in $X\mathbb{Z}H$, which gives $|G|m(|G| - \hat{G}) \in X\mathbb{Z}H$. Also

$$|G|\hat{G}(a_1 - 1) = |G|(x + \hat{G}(a_1 - 1)) - |G|x \in X\mathbb{Z}H,$$

whence $|G|^2m(a_1 - 1) = |G|m(|G| - \hat{G} + \hat{G})(a_1 - 1) \in X\mathbb{Z}H$. Therefore $|G|^2m\Delta H \subseteq X\mathbb{Z}H$, which completes the verification of (g2) and hence the proof of Proposition 6.1. $\square$

The following result is an immediate consequence of Theorem 4.6 and Proposition 6.1:

6.5. Theorem. *Let $H = H_1 * \dots * H_n$, where each H_i is of the form $G_i \times A_i$, A_i is a finitely generated nilpotent group with A_i/A_i' torsion-free and G_i is a finite group such that*

- (a) $d(G_i) = d(G_i/G_i')$ and
- (b) $d_{G_i}(\Delta G_i/p\Delta G_i) = d(G_i/G_i'G_i^p)$ for every $p \in \pi(G_i)$. Then

$$d_H(\Delta H) = \max_p \left\{ \sum_{i=1}^n d_{H_i}(\Delta H_i/p\Delta H_i) \right\}.$$

Theorem 2.2 is an immediate corollary of Theorem 6.5 provided we note that $\delta_{H_i}(\Delta H_i) = (1 - \delta_{A_i,1})d(A_i) + \delta_{A_i,1}$.

We now turn to relation modules. These are harder to deal with. Let G be a non-trivial finite group, let C be an infinite cyclic group, and let $H = G \times C$. Choose a finitely generated free presentation $R \hookrightarrow F \twoheadrightarrow G$ for G and form the finitely generated free presentation $S \hookrightarrow F * C \twoheadrightarrow H$ for H .

6.6. Proposition. $\overline{S}$ satisfies (g1) with $\delta_H(\overline{S}) = d(F)$.

Proof. Let $\alpha: F * C \twoheadrightarrow G * C$ denote the natural epimorphism induced by the surjection of F onto G and the identity on C , let $\beta: G * C \twoheadrightarrow G \times C$ denote the natural epimorphism induced by the identity on G and C , and let $\gamma: F * C \twoheadrightarrow G \times C$ denote the natural epimorphism induced by the surjection of F onto G and the identity on C . Set $R_0 = \ker \alpha$, $S = \ker \gamma$. Then we have a commutative diagram with exact rows

$$\begin{array}{ccccccccc} 1 & \longrightarrow & R_0 & \longrightarrow & F * C & \xrightarrow{\alpha} & G * C & \longrightarrow & 1 \\ & & \downarrow & & \parallel & & \downarrow \beta & & \\ 1 & \longrightarrow & S & \longrightarrow & F * C & \xrightarrow{\gamma} & G \times C & \longrightarrow & 1 \end{array}$$

where the map $R_0 \rightarrow S$ is the natural inclusion. Let $E = \ker \beta$ and write $C = \langle c \rangle$. By the snake lemma (or the map β can be identified with the natural map $(F * C)/R_0 \rightarrow (F * C)/S$), we see that $S/R_0 \cong E$. Also E is the free group on $\{[g, c^i] \mid 1 \neq g \in G, 0 \neq i \in \mathbb{Z}\}$, so $S = R_0 \rtimes E$, which gives the extension of $\mathbb{Z}H$ -modules

$$(6.7) \quad R_0/[R_0, S] \twoheadrightarrow \overline{S} \twoheadrightarrow E/E'.$$

Choose an inverse $t: G \rightarrow F$ to the free presentation $F \twoheadrightarrow G$ (so $t(G)$ is a transversal for R in F) and assume that $t(1) = 1$. Then $R_0 = *_w(w^{-1}Rw)$ (the free product of the subgroups $w^{-1}Rw$ where w runs through all those elements of $F * C$ that can be expressed as a word in $t(G)$ and C , and start with an element of C). Hence $\overline{R}_0 = \overline{R} \otimes_{\mathbb{Z}G} \mathbb{Z}[G * C]$ and then, because $R_0 \hookrightarrow S \twoheadrightarrow E$ is exact, $R_0/[R_0, S] = (\overline{R} \otimes_{\mathbb{Z}G} \mathbb{Z}[G * C]) \otimes_{\mathbb{Z}E} \mathbb{Z} \cong \overline{R} \otimes_{\mathbb{Z}G} \mathbb{Z}H \cong \overline{R} \otimes_{\mathbb{Z}} \mathbb{Z}C$; also $E/E' \simeq \Delta G \otimes_{\mathbb{Z}} \mathbb{Z}C$ as $\mathbb{Z}H$ -modules via $[g, c^i] \mapsto (g - 1) \otimes (c^i - 1)$ and $\Delta(C) \simeq \mathbb{Z}C$ via $c - 1 \mapsto 1$. Hence (6.7) becomes

$$(6.8) \quad \overline{R} \otimes_{\mathbb{Z}} \mathbb{Z}C \hookrightarrow \overline{S} \xrightarrow{\sigma} \Delta G \otimes_{\mathbb{Z}} \mathbb{Z}C.$$

Now $\tau: \Delta G \otimes_{\mathbb{Z}} \mathbb{Z}C \rightarrow \overline{S}$ defined by $(g - 1) \otimes c^i \mapsto \overline{c^{-i}[t(g), c]c^i} = \overline{[t(g), c]c^i}$ is a $\mathbb{Z}C$ -homomorphism, whence $\tau\hat{G}$ is a $\mathbb{Z}H$ -homomorphism. One checks that $(\tau\hat{G})\sigma = |G|\text{id}$.

So if q is prime to $|G|$, then (6.8) taken mod q is a split extension. The free presentation $R \hookrightarrow F \twoheadrightarrow G$ determines the relation sequence $\overline{R} \twoheadrightarrow \mathbb{Z}G^{d(F)} \twoheadrightarrow \Delta G$ and this splits modulo q . Thus $\overline{R}/\overline{R}q \oplus \Delta G/q\Delta G \cong (\mathbb{Z}G/q\mathbb{Z}G)^{d(F)}$ and therefore

$$\overline{S}/q\overline{S} \cong (\mathbb{Z}G/q\mathbb{Z}G)^{d(F)} \otimes_{\mathbb{Z}} \mathbb{Z}C/q\mathbb{Z}C \cong (\mathbb{Z}H/q\mathbb{Z}H)^{d(F)},$$

which gives $\delta_H(\overline{S}) = d(F)$.

Finally, let $p \in \pi(G)$ and write $E = F * C$. Then $\overline{S} \twoheadrightarrow SE'/E'$ is an H -homomorphism, SE'/E' is H -trivial and free abelian of rank $d(F)$, whence $\overline{S}/p\overline{S}$ has an H -trivial H -image of $\mathbb{F}_p$ -dimension $d(F)$. Thus $d_H(\overline{S}/p\overline{S}) \geq d(F)$. $\square$

Continuing with the notation of the last proof, take C -coinvariants of (6.8), giving

$$(6.9) \quad \overline{R} \hookrightarrow \overline{S} \otimes_{\mathbb{Z}C} \mathbb{Z} \rightarrow \Delta G,$$

which is exact because $H_1(C, \mathbb{Z}C) = 0$. The cocycle determining the extension (6.8) is $\psi: h \mapsto \tau h - \tau$. If $h = gc^i$, then $\psi(h) = \psi(g)$ because τ is a C -homomorphism. The cocycle determining (6.9) is $\psi_C := \psi \otimes_{\mathbb{Z}C} 1_{\mathbb{Z}}$.

Let $t(a)t(b) = t(ab)\rho(a, b)$ for $a, b \in G$. Then

$$\psi(g): (a-1) \otimes 1 \mapsto ((ag^{-1} - g^{-1}) \otimes 1)\tau g - ((a-1) \otimes 1)\tau = \bar{\rho}(a, g^{-1})g(1-c) \in \overline{S}\Delta C,$$

whence $\psi_C(g) = 0$. We conclude that (6.9) splits.

6.10. Lemma. *Assume that G is nilpotent and that there exists a relation prime of G that is also a ΔG -prime. Then $\overline{S}$ is a Swan module and every p that is both ΔG -prime and $\overline{R}$ -prime is an $\overline{S}$ -prime.*

Proof. Let p be an augmentation prime and also a relation prime for G . Then $d_G(\Delta G) = d(G/G'G^p)$ [9, Lemma 7.17(i)] and $d_G(\overline{R}) = d(R/[R, F]R^p)$ [9, Lemma 7.17(ii)]. Hence

$$d_G(\overline{R}) + d_G(\Delta G) = d(R/[R, F]R^p) + d(G/G'G^p).$$

Since (6.9) splits, there is an epimorphism $(\overline{S} \otimes_{\mathbb{Z}C} \mathbb{Z})/p(\overline{S} \otimes_{\mathbb{Z}C} \mathbb{Z}) \rightarrow R/[R, F]R^p \oplus G/G'G^p$, which shows

$$d(R/[R, F]R^p) + d(G/G'G^p) \leq d_G(\overline{S} \otimes_{\mathbb{Z}C} \mathbb{Z})/(\overline{S} \otimes_{\mathbb{Z}C} \mathbb{Z})p \leq d_H(\overline{S}/p\overline{S}) \leq d_H(\overline{S}).$$

But $d_H(\overline{S}) \leq d_G(\overline{R}) + d_G(\Delta G)$, by (6.8), so that we have achieved $d_H(\overline{S}) = d_H(\overline{S}/p\overline{S})$. $\square$

Unfortunately the nilpotence of G is not enough to ensure that the remaining hypothesis in Lemma 6.10 is true: cf. [9, Proposition 7.19]. But if G is cyclic (as it is from here on), then the set of all relation primes of G coincides with the set of all ΔG -primes and this set is $\pi(G)$.

6.11. Lemma. *Continue with the previous notation. If G is cyclic, then $\overline{S}$ satisfies (g2) for $\pi(G \times C, \overline{S}) = \pi(G)$.*

Proof. Let $n = |G|$, write $G = \langle a \rangle$, let $F = \langle x \rangle$ (so F is infinite cyclic), let $R = \langle x^n \rangle$, and let $x \mapsto a$ in the presentation $R \hookrightarrow F \rightarrow G$. Then S is the normal closure in $E = F * C$ of $\{x^n, [x, c]\}$, whence $[S, E]$ is generated mod E_3 (the third term $[[E, E], E]$ of the lower central series of E) by $[x^n, c] \equiv [x, c]^n \pmod{E_3}$. Thus $[S, E] \neq E'$, since E'/E_3 is infinite cyclic on $[x, c]E_3$. Therefore $d_H(\overline{S}) = 2$. Hence $d_H(\overline{S}/p\overline{S}) = 2$ for every $p \in \pi(G)$ (by 6.10).

The map $t: G \rightarrow F$ is here $t(a^i) = x^i$ for $0 \leq i < n$ and τ maps $(a-1) \otimes 1$ to $\overline{[x, c]}$. The calculation of $\psi(g)$ gives

$$\tau \hat{G}: (a-1) \otimes 1 \mapsto n\overline{[x, c]} + \left(\sum_{g \in G} \bar{\rho}(a, g^{-1})g \right)(1-c).$$

The term $\sum_{g \in G} \bar{\rho}(a, g^{-1})g = \overline{x^n}$. Set $z = n\overline{[x, c]} - \overline{x^n}c = ((a-1) \otimes 1)\tau \hat{G} - \overline{x^n}$. Since $\overline{S} = \{x^n, [x, c]\}\mathbb{Z}H$, we see that $\overline{S} = \{z, \overline{[x, c]}\}\mathbb{Z}H$. Now

$$z(n - \hat{G}c) = n^2\overline{[x, c]} - n\overline{x^n}c - \left(((a-1) \otimes 1)\tau \hat{G} - \overline{x^n} \right) \hat{G}c = n^2\overline{[x, c]},$$

which shows that $\overline{S} = z\mathbb{Z}H + q\overline{S}$ for any q prime to n ; and $\overline{S}/z\mathbb{Z}H$ has exponent dividing n^2 .

If π is a finite set of primes properly containing $\pi(G)$, then $\{z, \overline{[x, c]}\} \supseteq \{z\}$ is a nested π -family whose minimal element has the required properties for (g2). $\square$

Combining Proposition 6.6, Lemma 6.10 and Lemma 6.11 gives

6.12. Proposition. *If $F/R \simeq G$ is a minimal free presentation of the finite cyclic group G , C is an infinite cyclic group and $(F * C)/S \simeq G \times C$, then $\overline{S}$ is a good Swan module for $G \times C$ with $\pi(G \times C, \overline{S}) = \pi(G)$.*

We call the free presentation in Proposition 6.12 the *natural* minimal free presentation of $G \times C$. Proposition 6.12 and Theorem 4.6 combine to give

6.13. Theorem. *Let $H = H_1 * \cdots * H_n$, where each $H_i = G_i \times C_i$, G_i is finite cyclic and C_i is infinite cyclic. Take natural minimal free presentations $F_i/R_i \simeq H_i$ and let $F/R \simeq H$ be the resulting free presentation of H . Then*

$$d_H(\overline{R}) = \max_p \left\{ \sum_{i=1}^n d_{H_i}(\overline{R}_i/p\overline{R}_i) \right\}.$$

Finally in this section, we discuss [2, Proposition 2] (which is a generalisation of [2, Proposition 1]). Let n be a positive integer, let $\rho_n = \rho_n(x, t)$ be the word defined by $\rho_n(x, t) = (txt^{-1})x(txt^{-1})^{-1}x^{-n-1}$, and let $Q_n = \langle x, t \mid \rho_n = x^n = 1 \rangle$. Then [2, Lemma 3] shows that Q_n is isomorphic to the HNN-extension $(\mathbb{Z}/n\mathbb{Z} \times \mathbb{Z}/n\mathbb{Z}) *_{\phi}$, where ϕ maps the first factor isomorphically to the second. Let $R \hookrightarrow F \twoheadrightarrow Q_n$ be the presentation determined by the generators and relators for Q_n from above. Then, of course, the relation module $\overline{R}$ is generated as a $\mathbb{Z}Q_n$ -module by the images of ρ_n, x^n in $\overline{R}$. Define $q_n = (n+1)^n - 1$ and $c_n = nq_n$. Let $\bar{\rho}_n$ indicate the image of ρ_n in $\overline{R}$. Now [2, Lemma 4] proves that $[(txt^{-1})^n, x^n] = x^{c_n}$ in $\langle x, t \mid \rho_n \rangle$ (here $[a, b]$ denotes the commutator $aba^{-1}b^{-1}$). Furthermore since $[(txt^{-1})^n, x^n] \in R'$ and $\overline{R}/\bar{\rho}_n\mathbb{Z}Q_n$ is generated as a $\mathbb{Z}Q_n$ -module by the image of x^n in $\overline{R}/\bar{\rho}_n\mathbb{Z}Q_n$, we see that $\overline{R}/\bar{\rho}_n\mathbb{Z}Q_n$ is a group of exponent dividing q_n . We deduce that if p is a prime that does not divide q_n , then $\overline{R} = p\overline{R} + \bar{\rho}_n\mathbb{Z}Q_n$. We conclude that $\overline{R}$ is a good $\mathbb{Z}Q_n$ -module with $\delta_{Q_n}(\overline{R}) = 1$ and $\pi(Q_n, \overline{R}) = \pi(q_n)$, where $\pi(q_n)$ indicates the set of primes dividing q_n . Also if $n \geq 2$ and p is prime which divides n , it is not difficult to see that $d_G(\overline{R}/p\overline{R}) = 2$ and hence $\overline{R}$ is a Swan module. We can now apply Theorem 4.6 to obtain the following result.

6.14. Proposition. *Suppose r is a positive integer and $(q_{m_i}, q_{m_j}) = 1$ for $1 \leq i < j \leq r$. Assume that $m_i \geq 2$ for all i . If $\overline{R}$ is the relation module for the group $\Gamma := Q_{m_1} * \cdots * Q_{m_r}$ and presentation*

$$\langle x_{m_1}, t_{m_1}, \dots, x_{m_r}, t_{m_r} \mid \rho_{m_1}(x_{m_1}, t_{m_1}), \dots, \rho_{m_r}(x_{m_r}, t_{m_r}) \rangle,$$

then $d_{\Gamma}(\overline{R}) = r + 1$.

This result is essentially that of [2, Proposition 2], though there explicit generators for the relation module are given. Of course, one could obtain further results by combining the previous paragraph with Proposition 6.12 and Theorem 4.6.

7. APPENDIX

Throughout this section G is a finite group. Let M be a finitely generated $\mathbb{Z}G$ -module. As usual, $M_{(p)}$ denotes the localization of M at the prime p . To establish that M is a good $\mathbb{Z}G$ -module with $\pi(G, M) = \pi(G) \cup \pi(M)$ we must prove

7.1. Theorem. *Let π be a finite set of primes properly containing $\pi(G, M)$. Then there exists a nested π -family in M whose minimal element X satisfies*

$$d_G(M/(X\mathbb{Z}G + pM)) \leq 1$$

for every prime $p \notin \pi$.

We know that M has property (g1) (by Lemma 3.3) and $M/X\mathbb{Z}G$ is a finitely generated abelian group with $p(M/X\mathbb{Z}G) = M/X\mathbb{Z}G$ for $p \in \pi \setminus \{\pi(G) \cup \pi(M)\}$, whence $M/X\mathbb{Z}G$ is finite. Thus Theorem 7.1 supplies what is missing of (g2).

We recall some relevant facts. If R is a subring of $\mathbb{Q}$ containing $1/|G|$, then RG is a maximal order in $\mathbb{Q}G$, every central idempotent of $\mathbb{Q}G$ belongs to RG and every RG -lattice is RG -projective [4, respectively Proposition 27.1, Theorem 26.20, Theorem 26.12]. If U is an irreducible $\mathbb{Z}G$ -module such that $pU = 0$ for some $p \in \pi(G)'$, then U can be viewed as a $\mathbb{Z}_{(p)}G$ -module, whence there exists a unique primitive central idempotent e such that $Ue \neq 0$.

7.2. Lemma. *If V, W are projective $\mathbb{Z}_{(p)}G$ -lattices such that $\mathbb{Q}V \cong \mathbb{Q}W$, then $V \cong W$ [4, Theorem 32.1].*

7.3. Lemma. *Let M be a finite $\mathbb{Z}G$ -module such that $M_{(p)} = 0$ for all $p \in \pi(G)$. Then the composition factors of M can be arranged in any given order.*

This follows from [14, Lemma 9.3]. Here is another proof, shown to us by Steve Donkin.

Proof. It suffices to prove the result when M has prime-power order, so assume that $|M| = p^n$. Then $p \notin \pi(G)$ and M is a module over $\mathbb{Z}_{(p)}G$.

Let $M = M_0 \supset M_1 \supset \cdots \supset M_d = 0$ be a composition series and let $(U_1, \dots, U_d)$ be a (possibly new) sequence of the composition factors. We show by an induction on d that there exists a composition series of M producing factors in this order.

When $d = 1$ there is nothing to prove. Suppose $d = 2$. If $pM = 0$, then M is semi-simple and so we are done. If $pM \neq 0$, then pM must be simple and so is M/pM . Then $M \twoheadrightarrow pM$ has kernel pM , whence $M/pM \cong pM$ and again we are done.

Let $d > 2$ and $M/M_1 \cong U_r$. If $r = 1$, then induction gives a new series on M_1 that produces $(U_2, \dots, U_d)$, completing the proof. Suppose $r > 1$ and $U_1 \cong M_k/M_{k+1}$. Then $k > 0$ and by induction we can find a composition series of M_1 with U_1 at the top. By the case $d = 2$, there exists a composition series $M = M_0 \supset M'_1 \supset M'_2$ with factors (U_1, U_r) . Finally, use induction on M'_1 to obtain a composition series of M'_1 with factors $(U_2, \dots, U_d)$. $\square$

7.4. Lemma. *Let M be a finitely generated $\mathbb{Z}G$ -module, let p be a prime not in $\pi(G, M)$ and let U be an irreducible $\mathbb{Z}G$ -module such that $pU = 0$. Let e be the primitive central idempotent of $\mathbb{Q}G$ such that $Ue = U$ and assume that $(\mathbb{Q}M)e$ is not irreducible. If $\theta, \phi: M \twoheadrightarrow U$ are $\mathbb{Z}G$ -epimorphisms and q is an integer prime to p , then there exists a $\mathbb{Z}G$ -automorphism μ of M such that $\theta = \phi\mu$ and $\mu \equiv \text{id} \pmod{qM}$.*

Proof. Let M' be the torsion submodule of M . Since $p \nmid |M'|$, we see that $M_{(p)}$ is a $\mathbb{Z}_{(p)}G$ -lattice and $M_{(p)}e \neq 0$. If $(\mathbb{Q}M)e = V_1 \oplus V_2$ with $V_i \neq 0$, set $M_i = V_i \cap M_{(p)}e$ and then $M_1 \oplus M_2$ is a $\mathbb{Z}_{(p)}G$ -submodule of finite index in $M_{(p)}e$. Therefore $M_{(p)}e = M_1 \oplus M_2$ by Lemma 7.2 and there exist epimorphisms $\sigma_1: M_1 \twoheadrightarrow U$, $\sigma_2: M_2 \twoheadrightarrow U$.

Our proof follows that of [14, Proposition 9.5] but with some refinements. Given $\mathbb{Z}_{(p)}G$ -maps α_1, α_2 of M_1, M_2 to U , we shall write $\alpha_1 \oplus \alpha_2$ for the map $(x_1, x_2) \mapsto \alpha_1(x_1) + \alpha_2(x_2)$. Let θ_i be the map on M_i induced by θ . Clearly one of θ_1, θ_2 is an epimorphism. Let us suppose it is θ_2 .

Since M_1 is $\mathbb{Z}_{(p)}G$ -projective, there exists $\psi: M_1 \rightarrow M_2$ so that $\theta_2\psi = \theta_1 - \sigma_1$. Define $\tilde{\psi}: M_1 \oplus M_2 \rightarrow M_1 \oplus M_2$ by $(x_1, x_2) \mapsto (0, \psi(x_1))$. Then $\tilde{\psi}^2 = 0$. Since $M_{(p)}e$ is a direct summand of $M_{(p)}$, any endomorphism of $M_{(p)}e$ can be considered as one of $M_{(p)}$ by defining it to be zero on $M_{(p)}(1 - e)$. Then $\text{End}_{\mathbb{Z}_{(p)}G}(M_{(p)}) \cong \mathbb{Z}_{(p)} \otimes \text{End}_{\mathbb{Z}G}(M)$ shows $\tilde{\psi} = (1/r) \otimes \tau$ for a suitable integer r prime to p and $\tau \in \text{End}_{\mathbb{Z}G}(M)$.

As $|M'|, r, q$ are all prime to p , we may choose s so that $qr|M'|s \equiv 1 \pmod{p}$; then $t := |M'|s$ satisfies $tM' = 0$. Now $tM \cap M' = 0$ shows tM is a $\mathbb{Z}G$ -lattice and τ restricts to an endomorphism of tM . Since $(tM)_{(p)} \simeq M_{(p)}$ and $\tilde{\psi}^2 = 0$, we see that $\tau_{(p)}^2 = 0$, which gives $\tau^2(tM) = 0$. Define $\alpha = \text{id}_M + (qt)\tau$. This is a $\mathbb{Z}G$ -automorphism of M and we readily check that

$$(\sigma_1 \oplus \theta_2)\alpha_{(p)} = \theta_1 \oplus \theta_2.$$

A similar procedure leads to $\psi': M_2 \rightarrow M_1$ such that $\sigma_1\psi' = \sigma_2 - \theta_2$, $\tilde{\psi}': (x_1, x_2) \mapsto (\psi'(x_2), 0)$, $r'\tilde{\psi}' = \tau'_{(p)}$ for some r' coprime to p and some $\tau' \in \text{End}_{\mathbb{Z}G} M$. Then the $\mathbb{Z}G$ -automorphism $\alpha' = \text{id}_M + (qt)\tau'$ of M satisfies

$$(\sigma_1 \oplus \theta_2)\alpha'_{(p)} = \sigma_1 \oplus \sigma_2.$$

We conclude that the $\mathbb{Z}G$ -automorphism $\lambda = \alpha^{-1}\alpha'$ of M satisfies $(\theta_1 \oplus \theta_2)\lambda_{(p)} = \sigma_1 \oplus \sigma_2$. Similarly we can find an automorphism ν such that $(\phi_1 \oplus \phi_2)\nu_{(p)} = \sigma_1 \oplus \sigma_2$. Then $\mu = \nu\lambda^{-1}$ is an automorphism as required: for $\phi_{(p)}\mu_{(p)} = \theta_{(p)}$ implies $\phi\mu \equiv \theta \pmod{M'}$ (because $(p, |M'|) = 1$) whence $\phi\mu = \theta$. $\square$

7.5. Lemma. *Let L be a submodule of the $\mathbb{Z}G$ -lattice M such that M/L is finite of order prime to $|G|$. Then there exists a submodule N of M such that M/N is a finite cyclic $\mathbb{Z}G$ -module with the same composition factors as M/L .*

Proof. Since M/L is the direct sum of its primary parts, we may assume that M/L has p -power order for some prime p , so we may view M/L as a $\mathbb{Z}_{(p)}G$ -module. Let $V_1, \dots, V_n$ be the irreducible $\mathbb{Q}G$ -modules which appear in $\mathbb{Q}M$. Then we may choose a $\mathbb{Q}G$ -submodule V of $\mathbb{Q}M$ which is isomorphic to $\bigoplus_{i=1}^n V_i$. Set $A = V \cap M_{(p)}$. Since p is prime to $|G|$, all $\mathbb{Z}_{(p)}G$ -lattices are projective. Therefore we may write $M_{(p)} = A \oplus B$ for a suitable $\mathbb{Z}_{(p)}G$ -submodule B of $\mathbb{Q}M$, because $M_{(p)}/A$ is a $\mathbb{Z}_{(p)}G$ -lattice. Also we see from Lemma 7.2 that every submodule of A is cyclic, and if A_0 is a $\mathbb{Z}_{(p)}$ -submodule of A such that A/A_0 is finite, then every composition factor of M/L is a $\mathbb{Z}G$ -homomorphic image of A . Thus by induction on the length of the composition series of M/L , we see that there is a $\mathbb{Z}G$ -submodule C of A such that A/C has the same composition factors as M/L . Clearly A/C is a cyclic $\mathbb{Z}G$ -module. Now set $N = (C \oplus B) \cap M$. Then $M/N \cong A/C$ as $\mathbb{Z}G$ -modules and the result follows. $\square$

7.6. Lemma. *Let L, N be submodules of the finitely generated $\mathbb{Z}G$ -module M such that $M/L, M/N$ are finite with the same composition factors and $\pi(M/L) \cap \pi(G, M) = \emptyset$. Assume that if U is a composition factor of M/L with $pU = 0$ and attached to the primitive central idempotent e , then $(\mathbb{Q}M)e$ is not irreducible as $\mathbb{Q}G$ -module. Let q be any integer prime to $|M/L|$. Then there exists an isomorphism $\theta: L \simeq N$ such that $\theta(x) \equiv x \pmod{qM}$ for all $x \in L$.*

Proof. By Lemma 7.3 we may arrange the composition factors of M/L and M/N so that they appear in the same order; in other words, there exist chains of submodules $M = L_0 \supset L_1 \supset \cdots \supset L_d = L$ and $M = N_0 \supset N_1 \supset \cdots \supset N_d = N$ for some $d \geq 1$ such that, for each $i = 0, 1, \dots, d-1$, L_i/L_{i+1} is $\mathbb{Z}G$ -irreducible and $L_i/L_{i+1} \cong N_i/N_{i+1}$. We prove the result by induction on d .

By Lemma 7.4 there exists a $\mathbb{Z}G$ -automorphism μ of M such that μ induces $L_1 \simeq N_1$ and $\mu(x) \equiv x \pmod{qM}$ for all $x \in M$. By induction there exists an isomorphism $\nu: \mu L \simeq N$ such that $\nu(y) \equiv y \pmod{qN_1}$ for all $y \in \mu L$. Then $\nu\mu: L \simeq N$ and $\nu\mu(x) \equiv x \pmod{qM}$ for all $x \in L$. $\square$

Proof of Theorem 7.1. Choose a nested π -family $\mathcal{X}$ in M and let X be the minimal element. Let $e_1, \dots, e_n$ be the primitive central idempotents in $\mathbb{Q}G$ and define

$$\begin{aligned} I &= \{i \mid (\mathbb{Q}M)e_i \text{ is not irreducible}\}, \\ J &= \{i \mid (\mathbb{Q}M)e_i \text{ is irreducible}\}. \end{aligned}$$

Set $e = \sum_{i \in I} e_i$ and let $M_1/X\mathbb{Z}G$ be the π' -primary component of $M/X\mathbb{Z}G$. Define the $\mathbb{Z}G$ -submodule M_2 of M_1 containing $X\mathbb{Z}G$ by $M_2/X\mathbb{Z}G = (M_1/X\mathbb{Z}G)e$. Then each composition factor of $M_2/X\mathbb{Z}G$ is attached to a unique e_i with $i \in I$. The kernel of $M_2 \twoheadrightarrow M_2/X\mathbb{Z}G$ contains M'_2 , the torsion group of M_2 , because $\pi' \cap \pi(G) = \emptyset$.

By Lemma 7.5 there exists $N \supseteq M'_2$ such that M_2/N has the same composition factors as $M_2/X\mathbb{Z}G$ and M_2/N is a cyclic $\mathbb{Z}G$ -module. Let q be an integer prime to $|M_2/X\mathbb{Z}G|$. By Lemma 7.6 there exists an isomorphism $\theta: X\mathbb{Z}G \simeq N$ such that $x \equiv \theta(x) \pmod{qM}$ for all $x \in X$. Thus

$$X \equiv \theta(X) \pmod{pM}$$

for all $p \in \pi$. So if we replace the subset X of each member of the nested π -family $\mathcal{X}$ by $\theta(X) =: Y$, then we obtain a new nested π -family whose minimal element is Y . Observe $N = Y\mathbb{Z}G$ and $M/Y\mathbb{Z}G$ is finite because M/M_2 is finite. Also $(M_1/M_2)e = 0$, which makes $M_2/Y\mathbb{Z}G = (M_1/Y\mathbb{Z}G)e$.

We claim $M_1/Y\mathbb{Z}G$ is cyclic. Since we already know $(M_1/Y\mathbb{Z}G)e$ is cyclic, it remains to check $(M_1/Y\mathbb{Z}G)(1-e)$ is also cyclic. It suffices to do this for each primary component. If p occurs in the primary decomposition, then $(M_1)_{(p)}(1-e) = \sum_{i \in J} (M_1)_{(p)}e_i$, where $(\mathbb{Q}M)e_i$ for $i \in J$ are distinct irreducible modules, whence $(M_1)_{(p)}(1-e)$ is cyclic. Therefore so is $(M_1/Y\mathbb{Z}G)(1-e)$. Finally, if $p \in \pi'$, then $M/(pM + Y\mathbb{Z}G) = M_1/(pM + Y\mathbb{Z}G)$, which gives our result. $\square$

REFERENCES

- [1] George M. Bergman. Modules over coproducts of rings. *Trans. Amer. Math. Soc.*, 200:1–32, 1974.
- [2] Martin R. Bridson, Michael Tweedale, and Henry Wilton. Deficiency and abelianized deficiency of some virtually free groups. *Math. Proc. Camb. Phil. Soc.*, 143:257–264, 2007.

- [3] John Cossey, K. W. Gruenberg, and L. G. Kovács. The presentation rank of a direct product of finite groups. *J. Algebra*, 28:597–603, 1974.
- [4] Charles W. Curtis and Irving Reiner. *Methods of representation theory. Vol. I*. John Wiley & Sons Inc., New York, 1981. With applications to finite groups and orders, Pure and Applied Mathematics, A Wiley-Interscience Publication.
- [5] Francesca Dalla Volta and Andrea Lucchini. The smallest group with non-zero presentation rank. *J. Group Theory*, 2(2):147–155, 1999.
- [6] K. W. Gruenberg. Groups of non-zero presentation rank. In *Symposia Mathematica, Vol. XVII (Convegno sui Gruppi Infiniti, INDAM, Roma, 1973)*, pages 215–224. Academic Press, London, 1976.
- [7] K. W. Gruenberg. Free abelianised extensions of finite groups. In *Homological group theory (Proc. Sympos., Durham, 1977)*, volume 36 of *London Math. Soc. Lecture Note Ser.*, pages 71–104. Cambridge Univ. Press, Cambridge, 1979.
- [8] Karl W. Gruenberg. *Cohomological topics in group theory*. Lecture Notes in Mathematics, Vol. 143. Springer-Verlag, Berlin, 1970.
- [9] Karl W. Gruenberg. *Relation modules of finite groups*. American Mathematical Society, Providence, R.I., 1976. Conference Board of the Mathematical Sciences Regional Conference Series in Mathematics, No. 25.
- [10] Jens Harlander. Some aspects of efficiency. In *Groups—Korea '98 (Pusan)*, pages 165–180. de Gruyter, Berlin, 2000.
- [11] Cynthia Hog, Martin Lustig, and Wolfgang Metzler. Presentation classes, 3-manifolds and free products. In *Geometry and topology (College Park, Md., 1983/84)*, volume 1167 of *Lecture Notes in Math.*, pages 154–167. Springer, Berlin, 1985.
- [12] H. Jacobinski. Representation of orders over a Dedekind domain. In *Symposia Mathematica, Vol. XV (Convegno di Strutture in Corpi Algebrici, INDAM, Rome, 1973)*, pages 397–402. Academic Press, London, 1975.
- [13] F. E. A. Johnson. *Stable modules and the $D(2)$ -problem*, volume 301 of *London Mathematical Society Lecture Note Series*. Cambridge University Press, Cambridge, 2003.
- [14] Richard G. Swan. *K-theory of finite groups and orders*. Springer-Verlag, Berlin, 1970. Lecture Notes in Mathematics, Vol. 149.

SCHOOL OF MATHEMATICAL SCIENCES, QUEEN MARY, UNIVERSITY OF LONDON, MILE END ROAD, LONDON E1 4NS, ENGLAND

DEPARTMENT OF MATHEMATICS, VIRGINIA TECH, BLACKSBURG, VA 24061-0123, USA

E-mail address: linnell@math.vt.edu

URL: <http://www.math.vt.edu/people/linnell/>